

Research article

Open Access

Determination of Interleukin-6 and Tumor Necrosis Factor-alpha concentrations in Iranian-Khorasanian patients with preeclampsia

J Tavakkol Afshari*¹, N Ghomian^{†2}, A Shameli^{†2}, MT Shakeri^{†3},
MA Fahmidehkar^{†1}, E Mahajer¹, R Khoshnavaz⁴ and M Emadzadeh⁴

Address: ¹Immunogenetics department, Immunology Research Group, Bu-Ali Research Institute, Bu-Ali Sq., Mashhad University of Medical Sciences (MUMS), Mashhad, Iran, ²Obstetrics & Gynecology department, Imam Reza Hospital, Imam Reza Sq., Mashhad University of Medical Sciences (MUMS), Mashhad, Iran, ³Community medicine department, Medical school, Daneshgah St., Mashhad University of Medical Science (MUMS), Mashhad, Iran and ⁴Department of Cardiology, Ghaem Hospital, Ahmad Abad St., Mashhad University of Medical Sciences (MUMS), Mashhad, Iran

Email: J Tavakkol Afshari* - jtavakkol@yahoo.com; N Ghomian - N-ghomian@mum.ac.ir; A Shameli - Ahmad-shameli@mums.ac.ir; MT Shakeri - m_T_shakeri@yahoo.com; MA Fahmidehkar - mafahmidehkar@yahoo.com; E Mahajer - mmahajer@yahoo.com; R Khoshnavaz - Rasoulkhoshnavazmd@yahoo.com; M Emadzadeh - M-emadzadeh@mums.ac.ir

* Corresponding author †Equal contributors

Published: 01 November 2005

Received: 25 February 2005

BMC Pregnancy and Childbirth 2005, 5:14 doi:10.1186/1471-2393-5-14

Accepted: 01 November 2005

This article is available from: <http://www.biomedcentral.com/1471-2393/5/14>

© 2005 Afshari et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Our objective was to determine the role of Interleukin-6 (IL-6) and Tumor Necrosis Factor-alpha (TNF-alpha), markers of immune activation and endothelial dysfunction, in patients with preeclampsia.

Methods: Twenty four women with preeclampsia and eighteen antepartum normotensive pregnant women were recruited as controls. Serum levels of IL-6 and TNF-alpha were measured by enzyme-linked immunosorbent assay. We used independent-samples t test to assess the differences in the concentration of cytokines in preeclamptic patients and control subjects.

Results: IL-6 levels [mean (S.D.)] were significantly higher in preeclamptic women [5.8 (4.85) pg/ml] compared to normal pregnant women [3.01 (2.45) pg/ml] ($p = 0.02$). There was no significant change in concentration of TNF-alpha in preeclamptic women [53.8 (30.0) pg/ml] compared to normal pregnant women [51.9 (33.8) pg/ml] ($p > 0.1$).

Conclusion: The results of this study show that IL-6 as a pro-inflammatory cytokine is present in higher concentration in women with preeclampsia. The study was undertaken in women with established preeclampsia and it is not possible to determine whether the increased concentration of IL-6 is a cause or consequence of the disease. Furthermore, these findings suggest that serum TNF-alpha level is not associated with preeclampsia.

Background

Preeclampsia is a critically important disease of pregnancy, one of the major causes of fetal and maternal morbidity and mortality throughout the world. In spite of its importance for public health, the etiology of preeclampsia

has not yet been fully elucidated. Although the pathophysiology of preeclampsia is not understood completely, there is an interest in a possible link between inflammation and endothelial dysfunction [1] and between endothelial dysfunction and preeclampsia [2]. Cytokines

have been described to play a major role in the pathogenesis of preeclampsia. Recent studies have demonstrated that cytokines – mediators of inflammatory response–may cause endothelial dysfunction through different mechanisms such as oxidative stress [3] and endothelial cell damage [4].

IL-6 is a proinflammatory cytokine produced by mononuclear phagocytes, endothelial cells, fibroblasts and T cells and has many functions and effects [5]. IL-6 is involved in immune activation, vascular wall function and modulation of TNF- α production.

Recent studies have shown that amniotic fluid IL-6 is decreased in pregnancies complicated by preeclampsia and placental IL-6 production is decreased in these patients. These findings suggest a role for this cytokine in the pathophysiology of this disease. TNF- α is a potent modulator of immune and inflammatory responses that are produced by macrophages, lymphocytes and trophoblasts and contribute to the trophoblast growth and invasion. Its role in different pathologic conditions of pregnancy has been shown in recent studies. Increased amniotic fluid concentrations of TNF- α in patients with severe preeclampsia suggest a role for this cytokine in the pathophysiology of this disease. There are some studies in different populations that demonstrated the possible role of these cytokines in the pathophysiology of preeclampsia [6-10]. Therefore in the present study we measured their concentrations in serum of Khorasanian women (north-east of Iran) with preeclampsia, and compared them with concentrations found in normotensive control women.

Both IL-6 and TNF- α are expressed in adipose tissue [11,12] and *in vitro* release of TNF- α by adipocytes has been reported [13]. Among the known effects of these cytokines are inhibition of insulin signaling [14] and induction of both hypertriglyceridemia [15] and endothelial activation [16].

Methods

This study was approved by the ethical committee of the Mashhad University of Medical Sciences (MUMS) and was conducted at Imam-Reza Hospital and Bu-Ali Research Institute. It was a cross-sectional study and preeclamptic patients were selected from those admitted to Imam-Reza Hospital between February 2004 and December 2004. Written informed consent was obtained from participating women. Twenty four patients with preeclampsia at 26–41 weeks' gestation were enrolled in this study. Preeclampsia was diagnosed according to the clinical criteria defined by the "International Society for the study of Hypertension in Pregnancy" [17]. All preeclamptic patients had blood pressure of at least 140/90 mmHg and proteinuria greater than (+) as assessed by a

dipstick or 300 mg or greater in a 24-hour urine sample. None of the preeclamptic patients had underlying diseases such as chronic hypertension, diabetes mellitus, renal disease or urinary infection. None of these patients were in labor at the time of sampling. Some patients were prescribed hydralazine, dexamethazone or magnesium sulphate before sampling.

Eighteen antepartum normotensive healthy women at 26–39 weeks' gestation were included as control group. Maternal venous blood samples (5 ml) were collected from patient and control groups and were centrifuged at 3500 RPM for 10 minutes. Serum specimens were stored at -70°C in aliquots to avoid possible interference with assay results due to repeated freeze-thaw cycles.

Serum IL-6 and TNF- α concentrations were determined by commercially available high sensitivity indirect sandwich enzyme-linked immunosorbent assay (Bender MedSystems, Austria). Briefly, IL-6 or TNF- α present in the samples or standard binds to anti-IL-6 or anti-TNF- α monoclonal antibody adsorbed to the microwells. A biotin-conjugated monoclonal anti-IL-6 or anti-TNF- α antibody was added and binds to IL-6 or TNF- α captured by the first antibody. Following incubation unbound biotin-conjugated anti-IL-6 or anti-TNF- α is removed during a wash step. Streptavidin-HRP was added and binds to the biotin-conjugated anti-IL-6 or anti-TNF- α ; following incubation unbound Streptavidin-HRP was removed during a wash step, and substrate solution reactive with HRP was added to the wells. A colored product was formed in proportion to the amount of IL-6 or TNF- α present in the sample. The reaction was terminated by addition of acid and absorbance was measured at 450 nm. A standard curve was prepared from seven IL-6 and TNF- α standard dilutions and IL-6 or TNF- α sample concentrations determined. The detection limit for IL-6 and TNF- α were 1.4 pg/ml and 3.83 pg/ml respectively. The overall inter-assay coefficient of variation for IL-6 and TNF- α were 5.2% and 6.9% respectively. All serum IL-6 and TNF- α analyses were performed at the same time, in the same batch, and in duplicate according to manufacturer's instructions. IL-6 and TNF- α serum levels are given as normal distribution. Independent-samples *t* test was used to compare the mean levels between patient and control groups. *P*-value < 0.05 were considered statistically significant.

Results

A total of 24 women with preeclampsia and 18 women as controls were examined. The characteristics of women included in the study are presented in Table 1. There is no significance difference between two groups. Therefore the two groups were match for age, gestational age and gravity. IL-6 serum concentrations are depicted in Figure 1. IL-6 serum levels [mean (S.D.)] were significantly higher (*p*




Figure 1
Serum IL-6 concentration (pg/ml) measured by ELISA in normal pregnant (n-18) and preeclamptic (n = 24) women.




Figure 2
Serum TNF-alpha concentration (pg/ml) measured by ELISA in normal pregnant (n-18) and preeclamptic (n = 24) women.

= 0.02) in preeclamptic women [5.8 (4.85)] compared with normal pregnant women [3.01 (2.45)].

TNF- α serum concentrations are depicted in figure 2. There was no significant change in concentration of TNF- α serum levels in preeclamptic women [53.8 (30.0) pg/ml] compared to normal pregnant women [51.9 (33.8) pg/ml] ($p > 0.1$).

Discussion

Although the pathogenesis of preeclampsia is still unknown, immunologic and inflammatory causes may play an important role. IL-6 and other cytokines are important components of immune response, and therefore can participate in the immune aspects of the pathophysiology of this disease. Proinflammatory cytokines appear to be involved in cellular events that establish and maintain pregnancy [18]; however, their role has not yet been well defined.

Our data indicate that IL-6 concentrations were increased in the circulation of Iranian-Khorasanian preeclamptic patients compared with control women. This finding is consistent with reports by Greer et al in 1994 [6], Vince et al in 1995 [7], Kupferminc et al in 1996 [8], Munno et al in 1999 [9] and Teran et al in 2001 [10]. Endothelial dysfunction and increased endothelial permeability are the characteristics of the pathophysiology of preeclampsia. IL-6 may increase the permeability of endothelial cells [5] by changing the cell shape and rearrangement of intracellular actin fibers [19]. IL-6 can also reduce prostacyclin (PG I2)

synthesis by inhibiting the cyclooxygenase enzyme [20]. It can increase the tromboxane A2 to prostacyclin ratio, an abnormality that happens in preeclampsia. IL-6 can also stimulate platelet-derived growth factor, a process seen in preeclampsia [5]. Oxygen free radicals can induce synthesis of IL-6 by endothelium [21]. Oxygen free radicals are implicated in the pathogenesis of preeclampsia, because they can cause endothelial damage, which leads to reduction in nitric oxide synthesis and prostaglandin balance disturbance.

TNF- α is another pro-inflammatory cytokine that its contribution in the pathogenesis of preeclampsia has been suggested in recent studies [8,10]. In healthy pregnant women, TNF- α is thought to modulate the growth and invasion of trophoblasts in maternal spiral arteries [22]. TNF- α may contribute to abnormal placental invasion [23], endothelial cell damage [4] and oxidative stress [3]. TNF- α can stimulate IL-6 production [24], since IL-6 inhibits TNF- α release [25]. In contrast to IL-6, no increased serum concentration of TNF- α was found in preeclamptic patients compared to control pregnant women. Several investigators [26-28] have reported that serum concentrations of TNF- α were significantly higher in the first and second trimester among pregnant women who subsequently developed preeclampsia compared to those in the control group. This finding is consistent with reports by Greer et al in 1994 [6], Heyl et al in 1999 [28] and Ellis et al in 2001 [29]. Furthermore, the levels of IL-6 and TNF- α that we detected in the Khorasanian women studied were significantly higher than those reported for European and North American women [7,26,27]. Whether these differences are related to genetic (inflam-

Table 1:

	Normal pregnant (n = 18)	Preeclampsia (n = 24)	t and p
Age (years)	27.2 (5.8)	28.4 (4.9)	0.76, p > 0.1
Gestational age (weeks)	33.3 (3.7)	34.8 (4.3)	1.18, p > 0.1
Gravidity	2.3 (1.4)	2.2 (1.7)	0.65, p > 0.1

Data are presented as mean (S.D.).

matory response and L-arginine; NO pathway) and/or environmental factors (e.g., infection) remains to be determined. Several evidence links infection and inflammatory processes with preeclampsia [30,31]. The role of infection in the pathogenesis of preeclampsia is particularly relevant in developing countries, where the high incidence of chronic subclinical infection may contribute to the high incidence of preeclampsia.

Our findings support the hypothesis that immune activation is involved in preeclampsia and that IL-6 may participate in the abnormal immune response. This study was undertaken in women with established preeclampsia therefore, it cannot be determined whether the increase IL-6 was a cause or a consequence of the disease [7,10,26,27]. We cannot determine whether IL-6 is an active mediator in preeclampsia or a marker of immune activation, and it seems necessary to perform further studies, including longitudinal studies before the onset of preeclampsia, to elucidate the role of this cytokine in the pathogenesis of preeclampsia.

Conclusion

The results of this study show that IL-6 as a pro-inflammatory cytokine is present in higher concentration in women with preeclampsia. The study was undertaken in women with established preeclampsia and it is not possible to determine whether the increased concentration of IL-6 is a cause or consequence of the disease. Furthermore, these findings suggest that serum TNF- α level is not associated with preeclampsia.

Authors' contributions

JTA has participated in preparation and submission of the manuscript and given final approval of the version to be published. Also, conceived of the study, and participated in its design and coordination. NG participated in the diagnosis of patients with preeclampsia and in coordination of the study. AS participated in the diagnosis and selection of patients with preeclampsia. MTS participated in the design of the study and performed the statistical analysis. MAF carried out the immunoassays. MM participated in preparation of samples and performing ELISA for IL-6. RK: participated in drafting (English correction). ME participated in revising the manuscript critically for

important intellectual content. All authors read and approved the final manuscript.

Note

Table 1: Characteristics of women included in the study

Acknowledgements

This study was supported by the Vice Chancellor for Research, Mashhad University of Medical Sciences (MUMS), Mashhad, Iran.

References

- Vallance P, Collier J, Bhagat K: Infection, inflammation and infarction: does acute endothelial dysfunction provide a link? *Lancet* 1997, **349**:1391-1392.
- Roberts JM, Taylor RN, Musci TJ, Rodgers GM, Hubel CA, McLaughlin MK: **Preeclampsia: an endothelial cell disorder.** *Am J Obstet Gynecol* 1989, **161**:1200-1204.
- Stark JM: **Preeclampsia and cytokine induced oxidative stress.** *Br J Obstet Gynecol* 1993, **100**:105-109.
- Nawroth PP, Stern DM: **Modulation of endothelial cell hemostatic properties by tumor necrosis factor.** *J Exp Med* 1986, **163**:740-745.
- Akira S, Taniuchi T, Kishimoto T: **Interleukin-6 in biology and medicine.** *Adv Immunol* 1993, **54**:1-78.
- Greer IA, Lyall F, Perera T, Boswell F, Macara LM: **Increased concentrations of cytokines interleukin-6 and interleukin-1 receptor antagonist in plasma of women with preeclampsia: a mechanism for endothelial dysfunction?** *Obstet Gynecol* 1994, **84**:937-940.
- Vince GS, Starkey PM, Austgulen R, Kwiatkowski D, Redman CWG: **Interleukin-6, tumor necrosis factor and soluble tumor necrosis factor receptors in women with preeclampsia.** *Br J Obstet Gynecol* 1995, **102**:20-25.
- Kupferminc MJ, Peaceman AM, Aderka D, Wallach D, Socol ML: **Soluble tumor necrosis factor receptors and interleukin-6 levels in patients with severe preeclampsia.** *Obstet Gynecol* 1996, **88**:420-427.
- Munno I, Chiechi LM, Lacedra G, Berardesca C, Patimo C, Marcuccio L, Nardelli P, Loizzi P: **Evaluation of nonspecific immunity and plasma levels of interferon-gamma, interleukin-6 and tumor necrosis factor-alpha in preeclampsia.** *Immunopharmacol Immunotoxicol* 1999, **21**:551-564.
- Teran E, Escudero C, Moya W, Flores M, Vallance P, Lopez-Jaramillo P: **Elevated C-reactive protein and pro-inflammatory cytokines in Andean women with preeclampsia.** *Int J Gynecol Obstet* 2001, **75**:243-9.
- Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM: **Increased adipose tissue expression of tumor necrosis factor- α in human obesity and insulin resistance.** *J Clin Invest* 1995, **95**:2409-2415.
- Purohit A, Ghilchik MW, Duncan L, Wang DY, Singh A, Walker MM, Reed MJ: **Aromatase activity and interleukin-6 production by normal and malignant breast tissue.** *J Clin Endocrinol* 1995, **9**:61-72.
- Kern PA, Saghizadeh M, Ong JM, Bosch RJ, Deem R, Simsolo RB: **The expression of tumor necrosis factor in human adipose tissue; Regulation by obesity, weight loss, and relationship to lipoprotein lipase.** *J Clin Invest* 1995, **95**:2111-2119.
- Hotamisligil GS, Budavari A, Murray DL, Spiegelman BM: **Reduced tyrosine kinase activity of the insulin receptor in obesity-dia-**

- betes: central role of tumor necrosis factor- α .** *J Clin Invest* 1994, **94**:1543-1549.
15. Hardardóttir I, Grünfeld C, Feingold KR: **Effects of endotoxin and cytokines on lipid metabolism.** *Curr Opin Lipidol* 1994, **5**:207-215.
 16. van der Poll T, van Deventer SJH, Pasterkamp G, van Mourik JA, Büller HR, ten Cate JW: **Tumour necrosis factor induces von Willebrand factor release in healthy humans.** *Thromb Haemost* 1992, **67**:623-626.
 17. Davey DA, MacGillivray I: **The classification and definition of the hypertensive disorders of pregnancy.** *Am J Obstet Gynecol* 1988, **158**:892-898.
 18. Terranova PF, Hunter VJ, Roby KF, Hunt J: **Tumor necrosis factor-alpha in female reproductive tract.** *Proc Society Exp Biol Med* 1995, **209**:325-342.
 19. Maruo N, Morita I, Shirao M, Murota S: **IL-6 increases endothelial permeability in vitro.** *Endocrinology* 1992, **131**:710-714.
 20. Maruo N, Morita I, Ishizaki Y, Murota S: **Inhibitory effects of interleukin-6 on prostaglandin I2 production in cultured bovine vascular endothelial cells.** *Arch Biochem Biophys* 1992, **292**:600-604.
 21. Ala Y, Palluy O, Favero J, Bonne C, Modat G, Dornand J: **Hypoxia/reoxygenation stimulates endothelial cells to promote interleukin-1 and interleukin-6 production. Effects of free radical scavengers.** *Agents Actions* 1992, **37**:134-9.
 22. Kupfermink MJ, Peaceman AM, Wington TR, Tamura RK, Rehnberg KA, Socol ML: **Immunoreactive tumor necrosis factor- α is elevated in maternal plasma but undetected in amniotic fluid in the second trimester.** *Am J Obstet and Gynecol* 1994, **171**:976-979.
 23. Hunt JS, Soares MJ, Lei MG, Smith RN, Wheaton D, Atheron RA: **Products of lipopolysaccharide-activated macrophages (tumor necrosis factor, transforming growth factor) but not lipopolysaccharide modify DNA synthesis by rat trophoblast cells exhibiting the 80-kDa lipopolysaccharide-binding protein.** *J Immunol* 1989, **143**:1606-13.
 24. Shalaby MR, Waage A, Aarden L, Espevik T: **Endotoxin, tumor necrosis factor-alpha/cachectin and interleukin-1 induce interleukin-6 production in vivo.** *Clin Immunol Immunopathol* 1989, **53**:488-98.
 25. Aderka D, Le JM, Vilcek J: **IL-6 inhibits lipopolysaccharide-induced tumor necrosis factor production in cultured human monocytes, U937 cells, and in mice.** *J Immunol* 1989, **143**:3517-23.
 26. Williams M, Farrand A, Mittendorf R, Sorensen T, Zingheim R, Reily CO, King I, Zebelman A, Luthy D: **Maternal second trimester serum tumor necrosis factor-alpha-soluble receptor p55 (sTNFp55) and subsequent risk of preeclampsia.** *Am J Epidemiol* 1999, **149**:323-329.
 27. Sacks GP, Studena K, Sargent IL, Redman CWG: **Normal pregnancy and preeclampsia both produce inflammatory changes in peripheral blood leukocytes akin those of sepsis.** *Am J Obstet Gynecol* 1998, **179**:80-86.
 28. Heyl W, Handt S, Reister F, Gehlen J: **Elevated soluble adhesion molecules in women with preeclampsia. Do cytokines like tumor necrosis factor-alpha and interleukin-1beta cause endothelial activation.** *Eur J Obstet Gynecol Repord Biol* 1999, **86**:35-41.
 29. Ellis J, Wennerholm U, Bengtsson A, Lilja H: **Levels of dimethylarginines and cytokines in mild and severe preeclampsia.** *Acta Obstetrica Et Gynecologica Scandinavica* 2001, **80**:602-608.
 30. Hill JA, Devoe LD, Bryans CI Jr: **Frequency of asymptomatic bacteriuria in preeclampsia.** *Obstet Gynecol* 1986, **67**:529-532.
 31. Hsu CD, Witter FR: **Urogenital infection in preeclampsia.** *Int J Gynecol Obstet* 1995, **49**:271-275.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2393/5/14/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

