CORRECTION Open Access

Correction to: Early maternal weight gain as a risk factor for SGA in pregnancies with hyperemesis gravidarum: a 15-year hospital cohort study



Tale Meinich¹ and Jone Trovik^{1,2*}

Correction to: BMC Pregnancy and Childbirth (2020) 20:255 https://doi.org/10.1186/s12884-020-02947-3

Following publication of the original article [1], the authors identified an error in Fig. 1. The correct figure is given below.

The original article has been corrected.

Published online: 29 May 2020

Reference

 Meinich, Trovik. BMC Pregnancy Childbirth. 2020;20:255 https://doi.org/10. 1186/s12884-020-02947-3.

The original article can be found online at https://doi.org/10.1186/s12884-020-02947-3.

²Department of Obstetrics and Gynaecology, Haukeland University Hospital, Jonas Lies vei 72, 5053 Bergen, Norway



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*} Correspondence: jone.trovik@helse-bergen.no

¹Department of Clinical Science, University of Bergen, Jonas Lies vei 72, 5053 Bergen, Norway




Fig. 1 Weight gain patterns for 888 women hospitalized for hyperemesis gravidarum according to their prepregnancy body mass index (BMI) categories. Weight measured during 3-weeks interval during pregnancy until delivery. Comparisons by Kruskal-Wallis test